

SEAN P ROBINSON, MD Revolution Orthopedics

8120 Timberlake Way Suite 109 916 689 8441

Arthroscopic Ligament Reconstruction Postoperative instructions

Weight bearing

- You are touch down weight bearing on your operative leg with crutches
 - You will use crutches for 6 weeks due to your meniscus repair
- Keep your Brace on at all times you will be allowed to bend your knee to 90 degrees

Movement

- It is beneficial to change positions often after knee arthroscopy.
- o DO NOT STAY IN BED. We recommend at least 10 minutes of walking every hour.
- Work on patella mobilization

Sleep

- Keep your brace on during sleep or the first 2 weeks
- Keep your leg elevated.

• Ice

- Use bagged ice **as much as possible** when you get home at intervals of 30 minutes on and off. You should use ice for approximately two weeks. Do not use ice while you are sleeping.
 - Place the ice onto the knee over a thin layer of clothing or a towel, but never directly on the skin.

• Wound care, Dressings/Bandages

- Leave surgical bandage on and do not shower for 48 HOURS
- After 48 hours, remove bandages and gauze, but leave steri-strips (white tape) in place during showers.
- Its normal to see bloody soaked fluid on the bandages.
- After showers, leave the incisions open to air with steri-strips in place and gently pat dry. You may place BANDAIDS over the incisions
- o Do not apply lotions or ointments to the incisions.
- O not soak incisions in any pool/spa/bath water until 3 weeks after postoperatively or until your incisions are completely closed.

After surgery, call the clinic if you have....

- A fever higher than 101° (38.3° C).
- Changes in your incision: opening, drainage, redness.
- Numbness or tingling or loss of function of your leg.
 - Numbness around the incisions is normal.

- Increased pain or swelling in your calf.

 Nausea or other side effect not controlled by the medications you are given.

 CALL 911 IMMEDIATELY IF YOU HAVE CHEST PAIN OR SHORTNESS OF BREATH.